



VTE のトータルケア Web セミナーが

9月14日(水)19:30~20:30 に

第一三共㈱奈良営業所にて行われます。

当院からは

血管外科センター長 今井 崇裕 医師が

学術発表致しますので、ご紹介します。





Daiichi-Sankyo Oral AntiCoagulant Web Forum in Nara

VTEWebセミナー

VTEのトータルケア

「ここが変わった!静脈疾患治療」



座長

田村 大和先生

奈良県西和医療センター 心臓血管外科 部長

演者

今井 崇裕先生


康仁会 西の京病院 血管外科センター長

日時: 2016年 9月14日(水)

19:30~20:30

場所:第一三共株式会社 奈良営業所

当日は、ご施設名、ご芳名の記帳をお願い申し上げます。 ご記帳いただきましたご施設名、ご芳名は、医薬品の適正使用情報および医学・薬学に関する情報提供のために利用させて いただくことがございます。何卒、ご理解とご協力を賜りますようお願い申し上げます。



ASIAN SOCIETY FOR VASCULA SURGERY が

10月20日(木)~23日(日)に

シンガポールにて開催されます。

当院からは

血管外科センター長 今井崇裕 医師が

学術発表致しますので、ご紹介致します。



THE 17TH CONGRESS OF ASIAN SOCIETY FOR VASCULAR SURGERY

& 11TH ASIAN VENOUS FORUM



Selection criteria of endovenous ablation and stripping and its performing rate in our hospital

Takahiro Imai

Department of Vascular Surgery, Nishinokyo Hospital, Nara, Japan

Background and objectives: In 2011 Endovenous ablation for varicose vein was covered by national insurance and after that endovenous ablation are becoming increasingly popular in Japan. We started Endovenous ablation treatment using ELVeS laser 980nm in 2013 and we changed standard method to Radiofrequency Ablation using "ClosureFastTM" and stab avulsion in 2014. The ratio of Endovenous ablation for varicose veins gradually increases. In 2015, Stripping was 16.8% and ablation was 83.2%.

Methods: Endovenous ablation is a minimally invasive treatment and suitable for day surgery, but not applicable to all cases. We examined the factors affecting selection of operative method in our hospital.

Result: [Blood vessel diameter] According to the package insert provided with the ELVeS Laser device, blood vessel diameter must be less than 20mm. [Superficial varicose veins] Endovenous ablation is chosen, because we can secure distance between fiber and the skin surface by giving TLA. For young woman, we make a difficult choice considering a risk of pigmentation. [Growth of saphenofemoral junction tributaries] Endovenous Ablation is usually chosen. When the tributaries are not presumed to be completely occluded according to their locations on ultrasonography, stripping is chosen. [Patients on oral anticoagulants and antiplatelet drug] We perform Endovenous ablation with less bleeding during surgery. [Highly meandering blood vessel] Endovenous ablation is chosen because the guidewire is more likely to pass through the vein than stripper. [Varicose veins complicated by thrombophlebitis Stripping was chosen before, now Endovenous ablation is mostly chosen. [Recurrent case] We decide method considering the last operative method. [Operation invasion] In stripping, the surgical wound is small and the day surgery is possible, therefore operative method is not judged based on the patient wishes to have the day surgery or their cosmetic aspects.

Conclusions: Here we report examination results of the factors affecting selection of operative method.