
The 11th Japan-Korea Joint Meeting for Vascular Surgery





The 47th Annual Meeting of Japanese Society for Vascular Surgery



News

2019.01.17 Website has been opened.

Secretariat

Division of Vascular Surgery, Department of Surgery, Nagoya University Graduate School of Medicine 65 Tsurumai-cho, Showa-ku, Nagoya 466-8550, Japan
Phone: ±81-52-744-2224 / Fax: +81-52-744-2226 E-mail: jsys47@med_nagoya-u_ac_jp

Congress Secretariat of the 11th JKVS

c/o Congress Corporation

NARA Socks Project aiming at a spread of compression therapy and a revival of local industry

Takahiro Imai, MD, Department of Vascular Surgery, Nishinokyo Hospital, Nara, JAPAN

Abstract

BACKGROUND: In japan "Monozukuri" stands for the manufacturing industry and its spirituality and culture. It is not only making things but also Japanese traditional technique in pursuit of advanced production by the craftsmen. Koryo-cho with a population of 30 thousand called "Town of Socks" is the largest producer of socks traditionally here in Japan. These manufacturers are highly reputed for their technical skills and capabilities, and supply products to such overseas luxury brands. However, due to the large quantity of imported products to Japan, the production volume has dropped greatly from the peak period. And Japan is a country with many disasters because of island. It is necessary to enlightenment to prevention of thrombosis at the time of disaster to the people. I asked medical institutions, stocking manufacturers, and university students in local for cooperation. As a result, one elastic stocking was completed. The purpose of the activity is enlightenment of prevention of thrombosis and reconstruction of local industry. METHODS: The research period is from January 2017 to March. There are 30 objects (M14/F16, 34.5±9.3). We investigated changes in the circumference of the leg below the knee and improvements in the blood circulation of the legs before and after using the compression stockings with sonogram and investigated a compression pressure while wearing the stockings(Fig. 1).

RESULTS: The results revealed that the circumference of the leg below the knee showed a decreasing tendency after wearing the compression stockings. The peak velocity of the popliteal vein tended to increase 20 min after wearing the compression stockings (Fig.2). CONCLUSION: The compression stockings seemed to be effective for promoting venous return and preventing deep vein thrombosis of the lower limbs in healthy people. I am hoping to reduce the suffering caused by human thrombosis and revive the local industry.



| | | 0 | 20 | 40 |
|--------|-------------------------|-----------|-----------|-----------|
| Fig.2_ | Averaged PV (cm/second) | 4.65(1.9) | 5.62(2.1) | 5.51(2.3) |
| | VD (mm) | 8.65±1.1 | 8.48±1.2 | 8.39±1.2 |